

To appear in Ap. J.

A long-lasting quiescence phase of the eruptive variable V1118 Ori

D.Lorenzetti¹, S.Antoniucci¹, T.Giannini¹, A. Harutyunyan², A.A.Arkharov³,
V.M.Larionov^{3,4,5}, F.Cusano⁶, A.Di Paola¹, G.Li Causi¹, B.Nisini¹, R.Speziali¹, and
F.Vitali¹.

ABSTRACT

V1118 Ori is an eruptive variable belonging to the EXor class of Pre-Main Sequence stars whose episodic outbursts are attributed to disk accretion events. Since 2006, V1118 Ori is in the longest quiescence stage ever observed between two subsequent outbursts of its recent history. We present near-infrared photometry of V1118 Ori carried out during the last eight years, along with a complete spectroscopic coverage from 0.35 to 2.5 μm . A longterm sampling of V1118 Ori in quiescence has never been done, hence we can benefit from the current circumstance to determine the lowest values (i.e. the zeroes) of the parameters to be used as a reference for evaluating the physical changes typical of more active phases. A quiescence mass accretion rate between $1\text{--}3 \times 10^{-9} \text{ M}_{\odot} \text{ yr}^{-1}$ can be derived and the difference with previous determinations is discussed. From line emission and IR colors analysis a visual extinction of 1-2 mag is consistently derived, confirming that V1118 Ori (at least in quiescence) is a low-extinction T Tauri star with a bolometric luminosity of about $2.1 \text{ L}_{\odot}$. An anti-correlation

¹INAF - Osservatorio Astronomico di Roma, via Frascati 33, 00040 Monte Porzio, Italy, dario.lorenzetti, simone.antoniucci, teresa.giannini, andrea.dipaola, gianluca.licausi, brunella.nisini, roberto.speziali, fabrizio.vitali@oa-roma.inaf.it

²Fundación Galileo Galilei INAF, Telescopio Nazionale Galileo, 38700 Santa Cruz de la Palma, Tenerife, Spain, avet@tng.iac.es

³Central Astronomical Observatory of Pulkovo, Pulkovskoe shosse 65, 196140 St.Petersburg, Russia, arkadi@arharov.ru

⁴Astronomical Institute of St.Petersburg University, Russia, vlar2@yandex.ru

⁵Isaac Newton Institute of Chile, St.Petersburg branch

⁶INAF - Osservatorio Astronomico di Bologna, via Ranzani 1, 40127 Bologna, Italy, felice.cusano@oabo.inaf.it

exists between the equivalent width of the emission lines and the underlying continuum. We searched the literature for evaluating whether or not such a behaviour is a common feature of the whole class. The anti-correlation is clearly recognizable for all the available EXors in the optical range ($H\beta$ and $H\alpha$ lines), while it is not as much evident in the infrared ($Pa\beta$ and $Br\gamma$ lines). The observed anti-correlation supports the accretion-driven mechanism as the most likely to account for continuum variations.

Subject headings: Stars: pre-main sequence – variable – emission lines – Physical Data and Process: accretion disks – infrared: stars – individual: – individual: V1118 Ori

1. Introduction

After the accumulation of most of their final mass, young stellar objects (YSOs) of low-to-intermediate mass ($0.5\text{--}8\ M_{\odot}$) appear as pre-main sequence objects while the mass accretion process continues at a lower rate. In this phase the source is expected to accrete from its circumstellar disk: matter moves through the viscous disk and eventually falls onto the star surface following the magnetic interconnection lines (Shu et al. 1994). Observations show that the disk accretion process takes place through rapid and intermittent outbursts, usually detected at optical and near-IR wavelengths, which can be related to a sudden increase of the mass accretion rate by orders of magnitude (e.g. Hartmann & Kenyon 1985, Antonucci et al. 2008). An in-depth study of these accretion events is crucial to understand: *(i)* how the process eventually halts (thus determining the observed Initial Mass Function); *(ii)* how the bursts substantially alter the physical and chemical properties of the circumstellar disk, with major effects on the formation of proto-planetary systems; and *(iii)* the mechanism(s) at the origin of such outbursts.

Albeit a small and irregular photometric variations (typically 0.2-1 mag) caused by disk accretion variability is a defining feature of all classical T Tauri stars (CTTSs), several young sources display powerful outbursts of much larger intensity (up to 4-5 mag). Depending on different properties (burst duration, recurrence time between subsequent bursts, accretion rate, presence of absorption or emission lines), these objects are usually classified as FUors (Hartmann & Kenyon 1985) and EXors (Herbig 1989). These latter are more suitable for an observational monitoring, since their variations typically occur on shorter times: outbursts duration from some months to one year superposed to longer quiescence periods (years); moreover, they present accretion rates of the order of ($10^{-6}\text{--}10^{-7}\ M_{\odot}\ \text{yr}^{-1}$), and emission line spectra (e.g. Lorenzetti et al. 2009, Sicilia-Aguilar et al. 2012, Kóspál et al. 2011). The

reader is also referred to the excellent review by Audard (2014) who gives a complete view of the EXor properties.

Indeed, the nature of EXors is still very uncertain: no detailed analysis or modeling of their disk structure has been performed so far, so the mechanism responsible for the onset of the accretion outbursts is not known. Proposed scenarios involve thermal instability in the disk or gravitational instability triggered by a close companion (or planet). The latter alternative seems less probable, since in that case subsequent outbursts should occur with some periodicity instead of repeating at irregular time intervals.

The very uncertain picture of EXor events stems not only from the small number of known EXor objects (around two dozen, e.g. Lorenzetti et al. 2012; Audard et al. 2014), but especially from the lack of a proper multi-wavelength monitoring of their photometric and spectroscopic properties, which prevented the construction of a comprehensive database of information for these objects. For this reason, we started an observational program dubbed EXORCISM (EXOR OptiCal and Infrared Systematic Monitoring - Antonucci et al. 2013) that is intended to perform a photometric and spectral monitoring in the range 0.5-2.5 μm of about 20 objects identified as known eruptive variables (EXor) or candidates. In the framework of such a project, we present here optical and near-IR data of the typical source V1118 Ori collected during a quiescence phase in different dates from September 2006 to October 2014.

During its recent history V1118 Ori underwent five documented outbursts, each lasting a couple of years (1982-84, 1988-90, 1992-94, 1997-98, 2004-06). Account for the first four events is given in Parsamian et al. 1993, García García & Parsamian 2000, Herbig 2008 and references therein. For a complete view of the V1118 Ori properties shown during its last eruption, see Audard et al. 2005, 2010; Lorenzetti et al. 2006, 2007. Recently, Reipurth et al. (2007) discovered the binary nature of V1118 Ori, finding a close companion separated by 0.18 arcsec and about a factor 1.4 fainter in the $\text{H}\alpha$ filter (~ 0.4 mag).

Typically, outburst phases of EXors are monitored more frequently than quiescence ones, although these latter have exactly the same relevance (e.g. Sipos et al. 2009). In fact, only during these phases the lowest values (the zeroes) of the different parameters can be accurately evaluated so allowing one to compute the physical changes, once the outburst values are obtained. Moreover, because of their longer duration, the quiescence phases can be observationally traced more easily than outbursting ones.

Unfortunately, the quiescence status of V1118 Ori has never been thoroughly sampled, and in this paper we try to fill this gap. In Sect.2 optical and near-IR observations are presented; the results are analysed in Sect.3, while our concluding remarks are given in

Sect.4.

2. Observations

2.1. Near-IR imaging

Near-IR data were obtained at the 1.1m AZT-24 telescope located at Campo Imperatore (L'Aquila - Italy) equipped with the imager/spectrometer SWIRCAM (D'Alessio et al. 2000), which is based on a 256×256 HgCdTe PICNIC array. Photometry was performed with broad band filters J ($1.25 \mu\text{m}$), H ($1.65 \mu\text{m}$), and K ($2.20 \mu\text{m}$). All the observations were obtained by dithering the telescope around the pointed position. The raw imaging data were reduced by using standard procedures for bad pixel removal, flat fielding, and sky subtraction. Photometric data are listed in Table 1 while the derived light curves are depicted in the three panels of Figure 1 for the J, H, and K band, respectively. In the bottom panel (J band) we evidence two groups of activity phases: a long-term quiescence state (blue dots) and a short period of moderate activity (red dots). These states have been arbitrarily identified with a J-band magnitude greater (less) than 12.4, respectively. We will discuss these phases in deeper detail in Sect.3.1.

Near-IR data cover a period lasting eight years from September 2006 to October 2014; they have been collected without a systematic cadence and with no monitoring for long periods (of about one year), during which, however, no sign of outburst was reported (from both AAVSO circulars and Astronomer's Telegrams). Signs of modest activity (about 0.4 mag peak to peak) and short duration are superposed to such a long quiescence state (see Figure 1) that follows the last outburst, whose complete photometry is given elsewhere (Audard et al. 2010; Lorenzetti et al. 2006, 2007).

2.2. Optical spectroscopy

Optical spectra were taken on two different occasions (2014 March 25 and August 12, JD 2 456 741 and 2 456 881, respectively). The first was obtained with the 8.4m Large Binocular Telescope (LBT) using the Multi-Object Double Spectrograph (MODS - Pogge et al. 2010). The dual grating mode (Blue + Red channels) was used for a total integration time of 20 min to cover the $0.35 - 0.95 \mu\text{m}$ spectral range with a 0.6 arcsec slit (resolution ~ 2000). The second spectrum was obtained with the 3.6 m Telescopio Nazionale Galileo (TNG) using the Device Optimized for the LOw RESolution (DOLORES) instrument. The low resolution red (LR-R) grism was used for a total integration time of 30 min to cover the

0.50-0.95 μm spectral range with a resolution of ~ 700 . In both cases (LBT and TNG) images were bias and flat-field corrected using standard procedures. After removing sky background, the two-dimensional spectra were extracted and collapsed to one dimension. For both spectra, wavelength calibration was achieved through available lamp exposures, while spectral calibration was obtained from observations of spectro-photometric standards. The optical spectra of V1118 Ori are depicted in different colors (black for LBT, red for TNG) in Figure 2. These are the only quiescence spectra at a high level of sensitivity comparable with that of Herbig (2008) data. Both spectra show an extraordinary degree of repeatability concerning the continuum shape, whereas the derived line fluxes present variations of less than 20%. Such an occurrence might be the result of the extreme steadiness of this quiescence state, that, as such, can be considered very suitable for deriving the parameters of the object in its low state.

2.3. Near-IR spectroscopy

A low resolution spectrum ($\mathcal{R} \sim 500$, slit width 1 arcsec) was obtained on 2014 August 26 with NICS at TNG with two IR grisms IJ (0.90 - 1.45 μm) and HK (1.40 - 2.50 μm), in two subsequent exposures of 20 and 25 min, respectively. The standard ABB'A' mode was exploited within a long slit oriented at a position angle of 40° in order to minimize the flux of a close field star entering the slit. The spectral images were flat-fielded, sky-subtracted, and corrected for the optical distortion in both the spatial and spectral directions. Telluric features were removed by dividing the extracted spectra by that of a normalized telluric standard star, once corrected for its intrinsic spectral features. Wavelength and flux calibration were obtained from arc lamps and from our photometric data taken in the same period, respectively. In Figure 3 both portions (IJ and HK) of the resulting near-IR spectrum are given, labelling the relevant features identified.

3. Results and analysis

3.1. Near-IR photometry

Our images are not able to resolve the companion and no other high-angular resolution observations exist in other bands, hence we cannot precisely evaluate how the companion properties affect the quiescence parameters of V1118 Ori. Therefore, until this case will not be solved by future observations, we will not perform any (highly hypothetical) correction to the observed values derived for V1118 Ori, also considering that if a difference of 0.4 mag

maintains over the entire spectral range, such correction would be marginal.

In Figure 4 the near-IR two-colors plot is presented based on the photometric data given in Table 1. Data points are depicted with different colors (blue and red) to indicate the two different levels of activity evidenced in Figure 1 (see Sect.2.1). From the inspection of the two-colors plot, two considerations can be retrieved. First, all the data roughly cluster in the locus typical of T Tauri stars of a late spectral type and very low extinction. In particular, a value of $A_V \simeq 1\text{-}2$ mag can reasonably account for the data distribution. Second, a separation between the color of the quiescence phase (blue) and that of the moderate activity occurred during years 2008-2009 (points in red) is recognizable. In particular, we depicted with black filled dots the median values of the two distributions (considering 1σ error). The same trend (i.e. that different colors are associated even with modest continuum variations) has been found analysing also the light-curves of different EXors (Giannini et al. in preparation). Indeed, more significant decreases occurring in both the colors (i.e. a blueing effect) are commonly associated to the outburst phases of EXors (Lorenzetti et al. 2007, Audard et al. 2010; see e.g. Fig.1 of Lorenzetti et al. 2012). In this latter paper it is discussed how these variations cannot be accounted for simply by extinction, but rather by an additional thermal component of the emission. Such a circumstance seems to be confirmed also for the moderate activity presented here, since the red points are located fairly orthogonal to the extinction curve rather than along it.

Taking into account the lowest photometric values even detected, Audard et al. (2010) provided a plot of the quiescence Spectral Energy Distribution (SED) (their Figure 15-left) and an estimate of $2.0 L_\odot$ for the bolometric luminosity. By substituting their JHK photometry with our values taken on 2007 Mar 12, and by applying the on-line fitting procedure¹ (Robitaille et al. 2006, 2007), we obtain practically the same value of bolometric luminosity ($2.1 L_\odot$). Moreover, the output parameters (such as $T_\star = 4080$ K and $M_\star = 0.7 M_\odot$) are well compatible with the known physical parameters of V1118 Ori ($T_\star \simeq 3700$ K; $M_\star = 0.4 M_\odot$ - Hillenbrand 1997).

3.2. Optical and near-IR spectroscopy

The emission lines detected in the spectrum of the target are among those commonly observed in active young sources. A list of these lines is given in Table 2. The most prominent emission features are the H I recombination lines of the Balmer and Paschen series, which are commonly associated with flows of accreting gas (e.g. Calvet et al. 2000, 2004). Balmer

¹Available at <http://caravan.astro.wisc.edu/protostars/sedfitter.php>

lines are clearly visible up to H15 in the MODS spectrum, but no significant Balmer jump can be spotted at the end of the series, suggesting a fairly weak Balmer continuum. The spectrum shows also many permitted transitions of He I in the optical range and the He I line at $1.08 \mu\text{m}$, which is indicative of stellar winds (e.g. Edwards et al. 2006). Other permitted lines are those of Ca II (H and K line doublet at $0.39 \mu\text{m}$ and the triplet at $0.85\text{--}0.87 \mu\text{m}$) and the OI line at $0.845 \mu\text{m}$.

Weak forbidden emission lines of [O I] at $0.630 \mu\text{m}$ and [S II] at $0.636 \mu\text{m}$ (which are typical tracers of shocks driven by jets, see e.g. Giannini et al. 2008) are not detected in the spectrum of the source, although they are present as strong nebular contribution eliminated by the sky emission subtraction. This circumstance and the non-detection of [Fe II] (e.g. lines at $0.71\text{--}0.74$, 1.25 , and $1.64 \mu\text{m}$) and H_2 ($2.12 \mu\text{m}$) transitions indicate that, at least in the current quiescence state, there is no significant outflow activity from the source. This result is supported by the H_2 images of the Catalogue of Molecular Hydrogen Emission-Line Objects (MHOs - Davis et al. 2010), where some H_2 filamentary structures are present in the field, but unrelated with V1118 Ori.

From the observed emission lines we derived an estimate of the mass accretion rate ($\dot{M}_{acc}$) of the quiescent phase of V1118 Ori. For that we employed the set of empirical relationships that connect line and accretion luminosity (L_{acc}) derived by Alcalá et. al (2014) in a sample of young active T Tauri stars in Lupus. By adopting a distance of 400 pc, we computed the accretion luminosity from 24 different tracers, avoiding blended lines. The L_{acc} values thus inferred were then converted to mass accretion rates by using the relationship (e.g. Gullbring et al. 1998):

$$\dot{M}_{acc} = \frac{L_{acc} R_*}{GM_*} \left(1 - \frac{R_*}{R_{in}} \right)^{-1}, \quad (1)$$

where we assumed $M_* = 0.4 M_\odot$, $R_* = 1.29 R_\odot$ (Hillenbrand 1997, Stassun et al. 1999), and a typical inner radius $R_{in} = 5 R_*$. The $\dot{M}_{acc}$ values that we derive by adopting a visual extinction $A_V = 1$ are reported in Table 2. These are comprised in the interval 2×10^{-10} and 4×10^{-9} , with a median $\dot{M}_{acc}$ of about $1 \times 10^{-9} M_\odot \text{yr}^{-1}$ and a 1σ dispersion of $7 \times 10^{-10} M_\odot \text{yr}^{-1}$. By assuming $A_V = 2$ instead, we obtain a median accretion rate of about $3 \times 10^{-9} M_\odot \text{yr}^{-1}$ with a dispersion of $1.6 \times 10^{-9} M_\odot \text{yr}^{-1}$. The single accretion rates derived in both cases from all 24 lines are shown in Fig. 5.

We note that our $\dot{M}_{acc}$ estimates are much lower than the previously estimated quiescence $\dot{M}_{acc}$ of $2.5 \times 10^{-7} M_\odot \text{yr}^{-1}$ that was derived by Audard et al. (2010) from SED modelling; a further discussion on this aspect is given in Sect. 3.3.

3.3. Comparison with the outburst phase

We perform here a comparison between our inferred quiescence parameters and those derived during more active phases.

Bolometric luminosity L_{bol} - From the SED fitting, a value of $2.0 L_{\odot}$ is given (Audard et al. 2010 and confirmed in Sect.3.1) for the quiescent luminosity. Different estimates are given for the outburst luminosity: $25.4 L_{\odot}$ (Lorenzetti et al. 2006) and $7.4 L_{\odot}$ (Audard et al. 2010). These values refer to two different outbursts that likely exhibit different levels of brightness, nevertheless a substantial increase of the luminosity certainly occurs, related to the appearance of the spot onto the stellar surface and to the thermal disk contribution.

Visual extinction A_V - From the near IR two-colors plot (Figure 4) and from line emission analysis, A_V values of about 1-2 mag are consistently obtained. The extinction does not seem to change in the outburst during which Audard et al. (2010) derived, from X-rays column density, A_V values of $1.7^{+0.8}_{-0.6}$ mag or $1.4^{+0.6}_{-0.5}$ mag, for R_V equal to 3.1 or 5.5, respectively. A constant extinction suggests that changes of other physical parameters are not attributable to significant amounts of intervening dust (see also Audard et al. 2010; Lorenzetti et al. 2009).

Mass accretion rate $\dot{M}$ - In Sect. 3.2 $\dot{M}_{acc}$ values of 1 or $3 \times 10^{-9} M_{\odot} \text{yr}^{-1}$ were derived, depending on the assumed A_V , 1 or 2 mag, respectively. This value is two orders of magnitude lower than the previously estimated quiescence $\dot{M}_{acc}$ of $2.5 \times 10^{-7} M_{\odot} \text{yr}^{-1}$ derived by Audard et al. (2010) from SED modelling. This discrepancy is likely related to the different methods employed for the computation of the accretion rate. Accretion estimates from the emission lines are to be regarded as more reliable, since the lines are believed to trace the accretion columns or the strong accretion-related winds from the object. Moreover, the empirical accretion luminosity-line luminosity relationships, which are widely used as a proxy for deriving $\dot{M}_{acc}$, are directly calibrated from measurements of the UV excess emission from the accretion shock (e.g. Alcalá et al. 2014, Muzerolle et al. 1998). Finally, we note that the $\dot{M}_{acc}$ values we find for the quiescence of V1118 Ori are completely consistent with the accretion rates measured in most T Tauri objects (e.g. Natta et al. 2006, Alcalá et al. 2008, Biazzo et al. 2012, Antonucci et al. 2011, 2014).

For a comparison, we applied the emission line method to compute $\dot{M}_{acc}$ to our 2006 data (note that at that time the relationships for deriving $\dot{M}_{acc}$ from line fluxes were not available yet). From three bright Paschen lines ($\text{Pa}\beta$, $\text{Pa}\gamma$, and $\text{Pa}\delta$) we get $\dot{M}_{acc}$ value $\sim 1.0 \times 10^{-7} M_{\odot} \text{yr}^{-1}$ namely around 2 orders of magnitude higher than the quiescence value. Notably, this value is lower not only than the outburst value of Audard et al. (2010) of $1.0 \times 10^{-6} M_{\odot} \text{yr}^{-1}$ but even than their quiescence value of $2.5 \times 10^{-7} M_{\odot} \text{yr}^{-1}$. While the

difference between the two outburst values could be ascribed to the different outburst phases sampled in Lorenzetti et al. (2006) and in Audard et al. (2010), the inconsistency between our outburst value and the quiescence one by Audard et al. can only be a result of the different computation methods. Finally, we note that our outburst determination of $\dot{M}_{acc} / \dot{M}_{loss}$ is within the range (roughly 2-20) predicted by jet launching models (Shu et al. 2000, Königl & Pudritz 2000, Ferreira, Dougados & Cabrit, 2006), if we take the value of $\dot{M}_{loss} = 4.0 \times 10^{-8} M_{\odot} \text{ yr}^{-1}$ obtained from the HI recombination lines.

Emission lines - As listed in Table 2 the optical-IR quiescence spectrum of V1118 Ori is dominated by HI and He I recombination lines. Very few and faint metallic lines are also present. Such a situation is the same reported by Herbig (2008), who observed the source twice, one in outburst and one in a fading phase. While in the former phase he noticed the prominence of H I, He I, Ca II, Fe I, Fe II, and other neutral and ionized metals, when the star was much fainter the spectrum had changed radically: only HI and He I emission lines remained strong. Also our near-IR spectrum, taken during the last outburst (Lorenzetti et al. 2006) showed metallic (e.g. Na I) and CO emission features, now completely disappeared, even if we observed at higher sensitivity. In particular, CO emission is usually associated with large amounts of warm gas ($T \sim 3000$ K) in the inner regions of the circumstellar disc, which is indicative of an active phase of accretion (Najita et al. 1996; Lorenzetti et al. 2009; Kóspál et al. 2011). During a similar quiescence phase, Herbig (2008) observed also TiO absorption bands and the Li I 6707 absorption line. While TiO bands are clearly observed in our spectrum, Li I is absent, possibly because of a superposition of emission and absorption, both recurrent on different activity stages of V1118 Ori (Herbig 2008). However, we should have been able to detect a faint Li I feature similar to that observed by Herbig at just a 3σ level ($\simeq 7 \times 10^{-17} \text{ erg s}^{-1} \text{ cm}^{-2}$) over the continuum. In our quiescence spectrum Ca II emission lines are also present, but no comparison can be made with the Herbig (2008) spectrum that did not cover this spectral range. However, the presence of Ca II lines cannot be considered a peculiarity, as they are quite ubiquitous in T Tauri systems.

3.4. Equivalent widths and continuum

In the general context of the debate on whether the EXor events are accretion or extinction driven, it is worthwhile to analyse the relationship between lines and continuum emission. In particular, we aim at comparing the line emission behaviour in phases characterized by a different level of activity, i.e. how the EW varies with the increasing underlying continuum.

First of all we notice that the only member of the EXor class analyzed in this respect

(PVCep) shows a clear correlation (typical regression coefficient ≥ 0.9) between line and continuum flux (Lorenzetti et al. 2013). Since HI-recombination lines are considered as good tracers of the accretion rate (e.g. Muzerolle et al. 1998) the observed correlation supports the idea that accretion-driven mechanism is the most likely to account for continuum variations. Notice that accretion or mass loss processes are both compatible with the observed behaviour, since their ratio $\dot{M}_{acc} / \dot{M}_{loss}$ is roughly in the range 2-20. Although line and continuum fluxes are correlated, we can provide a more quantitative information by directly comparing the line EW and the continuum itself.

Literature data concerning the EW variations of prominent emission lines and those of their underlying continua are taken starting from the list of EXors and candidates (Lorenzetti et al. 2012, their Table 1). We report in Tables 3 and 4 only the data that refer to simultaneous observations of HI recombination lines $H\beta$ and $H\alpha$ and the continuum in the bands B and R, respectively. Providing that intra-day or day time-scale variations are usually expected to be modest, in few occasions, the values of EW and continuum have been assigned to the same date (within a maximum distance of 3-4 days), even if not strictly simultaneous. In Figure 6 the results of our literature search are depicted. The same exercise was done for the near-IR recombination lines $Pa\beta$, $Br\gamma$ and the relative photometry in the J, K bands, respectively. For the sake of compactness, near-IR data are not listed in a Table, but are also presented in the same Figure 6 and discussed below.

The anti-correlation, evident for V1118, occurs also for the majority of other sources when the optical lines ($H\beta$ and $H\alpha$) are considered, while for the near-IR lines ($Pa\beta$ and $Br\gamma$) the anti-correlation is not as much evident, being ascertainable only for about 50% of the presented cases. However, if we take into account only the most significant variations, namely those with $\Delta EW \geq 100\%$, or $\Delta mag \geq 2$, the total percentage (considering both optical and near-IR lines) of anti-correlations increases to about 80%. In conclusion, the existence of an anti-correlation can be reasonably confirmed as a general property of the EXor class of objects: a result already supported by previous investigations on individual objects (e.g. Cohen et al. 1981, Magakian & Movsessian 2001, Acosta-Pulido et al. 2007).

The anti-correlation indicates that the continuum presents larger (likely faster) variations than those of the lines, which means that continuum and lines do not obey to a common mechanism of heating and cooling. Such a circumstance tends to rule out the variable extinction as cause of the observed variability, since in that case a constant value of EW should be expected for any continuum fluctuation. Even a selective obscuration can be ruled out: dust should be located at the dust condensation zone, very close to the star and, as such, it should obscure the stellar photosphere more than the accretion columns and the wind regions.

4. Concluding remarks

We have presented the results of a monitoring of the EXor system V1118 Ori. Data concerning both IR photometry and optical/near IR spectroscopy cover a long lasting period of 8 years (2006-2014). This period starts from the last outburst and is the longest state of quiescence ever monitored. Hence it is very suitable for deriving reliable parameters typical of this status, that can be used as a reference to evaluate physical changes occurring during more active phases.

- Near IR colors of V1118 Ori cluster in a locus typical of a late type T Tauri star embedded in a low-extinction ($A_V \simeq 1\text{-}2$ mag) environment. A trend of the colors to become bluer when the source flux increases, can be recognized even in presence of modest ($\simeq 0.4$ mag) variations.
- From model fitting, stellar parameters ($T_\star = 3900$ K; $M_\star = 0.7 M_\odot$) well compatible with the known physical parameters of V1118 Ori are derived and the value of $2.0 L_\odot$ for the bolometric luminosity in quiescence is confirmed.
- Optical/near IR spectra are dominated by both H I recombination lines and He I permitted transitions. TiO absorption bands are also identified, without any sign of forbidden transitions, indicating that (at least in the current quiescence state) there is no significant outflow activity from the source.
- Emission line fluxes are consistently exploited to derive a mass accretion rate ($\dot{M}$) of $1\text{--}3 \times 10^{-9} M_\odot \text{ yr}^{-1}$ compatible with typical values observed in T Tauri stars. This value is substantially different from that previously derived using a different method. This discrepancy is likely related to the different methods employed for the computation of the accretion rate. Accretion estimates from the emission lines are to be regarded as more reliable, since the lines are believed to trace the accretion columns or the strong accretion-related winds from the object.
- An (anti-)correlation exists between the equivalent width of the emission lines and the underlying continuum. A search in the literature confirms that the behaviour of V1118 Ori is a common feature of the whole class and supports the accretion-driven mechanism as the most likely to account for continuum variations.

In Table 5 we summarize the results presented above: this table can be a useful reference for delineating the V1118 Ori properties at quiescence.

5. Acknowledgements

Based on observations made with different instrument. [1] the Italian Telescope Galileo (TNG) operated on the island of La Palma by the Fundación Galileo Galilei of the INAF (Istituto Nazionale di Astrofisica) at the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofisica de Canarias; [2] the Large Binocular Telescope (LBT). The LBT is an international collaboration among institutions in the United States, Italy and Germany. LBT Corporation partners are: The University of Arizona on behalf of the Arizona university system; Istituto Nazionale di Astrofisica, Italy; LBT Beteiligungsgesellschaft, Germany, representing the Max-Planck Society, the Astrophysical Institute Potsdam, and Heidelberg University; The Ohio State University, and The Research Corporation, on behalf of The University of Notre Dame, University of Minnesota and University of Virginia. [3] the AZT-24 IR Telescope at Campo Imperatore (L'Aquila - Italy) operated under the responsibility of the INAF-Osservatorio Astronomico di Roma (OAR).

REFERENCES

- Acosta-Pulido, J.A. et al. 2007, AJ, 133, 2020
- Alcalá, J. M., et al. 2008, ApJ, 676, 427
- Alcalá, J. M., Natta, A., Manara, C. F., et al. 2014, A&A, 561, 2
- Antoniucci, S. et al. 2011, A&A, 534, 32
- Antoniucci, S. et al. 2013, *Protostars and Planets VI*, on-line posters
<http://www.mpia-hd.mpg.de/homes/ppvi/posters/2B055.html>
- Antoniucci, S., García Lopez, R., Nisini, B., Caratti o Garatti, A., Giannini, T., & Lorenzetti, D. 2014 arXiv 1410.0181
- Antoniucci, S., Nisini, B., Giannini, T., & Lorenzetti, D. 2008, A&A 479, 503
- Aspin, C. 2011a, AJ, 141, 196
- Aspin, C. 2011b, AJ, 142, 135
- Aspin, C. et al. 2009, ApJ, 692, L67
- Aspin, C., & Reipurth, B. 2009, AJ, 138, 1137
- Audard, M. et al. 2014 Protostars and Planets VI - University of Arizona Press; eds. Beuther, H., Klessen, R., Dullemond, C., Henning, Th., in press
- Audard, M., Güdel, M., Skinner, S.L. et al. 2005, ApJ Lett. 635, L81
- Audard, M., Stringfellow, G.S., Güdel, M. et al. 2010, A&A 511, 63
- Biazzo, K., Alcalá, J. M., Covino, E., Frasca, A., Getman, F., & Spezzi, L. 2012, A&A, 547, 104
- Calvet, N., Hartmann, L., & Strom, S.E. 2000, Protostars and Planets IV - University of Arizona Press; eds V.Mannings, A.P.Boss, S.S Russell, p.377
- Calvet, N., Muzerolle, J., Briceno, C., et al. 2004, AJ, 128, 1294
- Cardelli, J.A., Clayton, G.C. & Mathis, J.S. 1989, ApJ, 345, 245
- Cohen, M., Kuhl, L.V., Harlan, E.A., & Spinrad, H. 1981, ApJ, 245, 920

- D'Alessio, F., et al. 2000, Proc. of the SPIE Symp. on *Astronomical Telescopes and Instrumentation*, eds. M. Iye & A.F.M. Moorwood, 4008, 748
- Davis, C.J., Gell, R., Khanzadyan, T., Smith, M.D., & Jenness, T. 2010, A&A, 511, 24
- Edwards, S., Fischer, W., Hillenbrand, L., & Kwan, J. 2006, ApJ, 646, 319
- Ferreira, J., Dougados, C., & Cabrit, S. 2006, A&A, 453, 785
- García García, J. & Parsamian, E.S. 2000 *Information Bulletin of Variable Stars* n.4925
- Giannini, T., Calzoletti, L., Nisini, B., et al. 2008, A&A, 481, 123
- Gullbring, E., Hartmann, L., Bričeno, C., & Calvet, N. 1998, ApJ, 492, 323
- Hartmann, L., & Kenyon, S. 1985, ApJ, 299, 462
- Herbig, G.H. 1989, Proc. of the ESO Workshop on *Low Mass Star Formation and Pre-Main Sequence Objects*, ed. B. Reipurth, p.233
- Herbig, G.H. 2008, AJ, 135, 637
- Herbig, G.H., Aspin, C., Gilmore, A.C., Imhoff, C.L., Jones, A.F. 2001, PASP, 113, 1547
- Hillenbrand, L. A. 1997, AJ, 113, 1733
- Königl, A. & Pudritz, R.E. 2000 Protostars and Planets IV - University of Arizona Press; eds V.Mannings, A.P.Boss, S.S Russell, p.759
- Kóspál, Á. et al. 2011, A&A, 527, A133
- Kun, M. et al. 2011, ApJL, 733, L8
- Lorenzetti, D., Antonucci, S., Giannini, T., Di Paola, A., Arkharov, A.A., & Larionov, V.M. 2013, Ap&SS, 343, 535
- Lorenzetti, D. et al. 2006, A&A, 453, 579
- Lorenzetti, D. et al. 2012, ApJ, 749, 188
- Lorenzetti, D., Giannini, T., Larionov, V.M., Kopatskaya, E., Arkharov, A.A., De Luca, M., & Di Paola, A. 2007, ApJ, 665, 1193
- Lorenzetti, D., Larionov, V.M., Giannini, T., Arkharov, A.A., Antonucci, S., Nisini, B., & Di Paola, A. 2009, ApJ, 693, 1056

- Magakian, T., & Movsesian, T.A. 2001, *Astrophysics* 44, 419
- Meyer, M.R., Calvet, N. & Hillenbrand, L.A. 1997, *AJ*, 114, 288
- Muzerolle, J., Hartmann, L., & Calvet, N. 1998, *AJ*, 116, 2965
- Natta, A., Testi, L., & Randich, S. 2006, *A&A*, 452, 245
- Ojha, D.K. et al. 2006 *MNRAS* 368, 825
- Parsamian, E.S., Gasparian, K.G., Oganian, G.B. & Melkonian, A.S. 1996 *Astrophysics* 39, 201
- Parsamian, E.S., Ibragimov, M.A., Oganian, G.B. & Gasparian, K.G. 1993 *Astrophysics* 36, 23
- Parsamian, E.S., Mujica, R. & Corral, L. 2002 *Astrophysics* 45, 393
- Pogge, R.W., Atwood, B., Brewer, D.F., et al. 2010, *Proc. SPIE*, 7735, 9
- Reipurth, B., Guimarães, M. M., Connelley, M. S., & Bally, J. 2007, *AJ*, 134, 2272
- Robitaille, T.P., Withney, B., Indebetouw, R., Wood, K., & Denzmore, P. 2006, *ApJS*, 167, 256
- Robitaille, T.P., Withney, B., Indebetouw, R., & Wood, K. 2007, *ApJS*, 169, 328
- Shu, F.H., Najita, J.R., Ostriker, E., Wilkin, F., Ruden, S., & Lizano, S. 1994, *ApJ*, 429, 781
- Shu, F.H., Najita, J.R., Shang, -H. & Li, Z.-Y. 2000, *Protostars and Planets IV* - University of Arizona Press; eds V.Mannings, A.P.Boss, S.S Russell, p.789
- Sicilia-Aguilar, A. et al. 2008, *ApJ*, 673, 382
- Sicilia-Aguilar, A. et al. 2012, *A&A*, 507, 93
- Sipos, N. et al. 2009, *A&A*, 507, 881
- Stassun, K. G., Mathieu, R. D., Mazeh, T., & Vrba, F. J. 1999, *AJ*, 117, 2941




Fig. 1.— JHK light-curves of V1118 Ori. In the lower panel (J band) data points corresponding to the highest flux levels are depicted in red, while in blue those typical of a more quiescent state.




Fig. 2.— Optical spectra of V1118 Ori, taken with LBT/MODS (black) and TNG/DOLORES (red). Main emission features are indicated.




Fig. 3.— Near-IR spectrum of V1118 Ori, taken with TNG/NICS IJ (red) and HK (blue) grisms; spectral segments that were heavily corrupted by telluric absorptions were removed. Main emission features are indicated.




Fig. 4.— Near-IR two colors plot ($[J-H]$ vs. $[H-K]$) of the quiescence period of V1118 Ori. Data points are those given in Table 1. A separation between the color of the quiescence phase (blue - the same as in Figure 1) and that of the moderate activity (red) is recognizable. Black solid circles indicate the median values of the two distribution (considering 1σ error). The solid line marks the unreddened main sequence, whereas the dashed one is the locus of the T Tauri stars (Meyer et al. 1997). Black dotted line represents the reddening law by Cardelli et al. (1989), where different values of A_V are indicated by open squares.




Fig. 5.— Mass accretion rate determinations computed from the flux of the indicated tracers using Alcalá et al. (2014) relationships, by assuming a visual extinction $A_V = 1$ (blue circles) and $A_V = 2$ (red squares). The median $\dot{M}_{acc}$ values are marked for both cases with a solid line.




Fig. 6.— EW of optical (upper panels) and IR (lower panels) HI recombination lines as a function of the underlying continuum expressed in magnitudes. To avoid a possible confusion, the EW values are labelled (in the y-axis) with a positive sign, although they are conventionally negative for emission lines, as reported in Tables 3 and 4.

Table 1. Near-IR photometry of V1118 Ori.

Date (yy/mm/dd)	JD +2400000	J	H (mag)	K	Date (yy/mm/dd)	JD +2400000	J	H (mag)	K
06 Sep 06	53984.6	12.63	11.61	10.96	07 Nov 05	54410.6	12.45	11.44	10.66
06 Sep 07	53985.6	12.62	11.66	10.97	08 Mar 29	54555.3	12.67	11.70	11.03
06 Sep 11	53989.6	12.66	11.66	10.98	08 Sep 06	54715.6	12.47	11.46	10.70
06 Sep 22	54000.6	12.63	11.57	10.82	08 Sep 29	54738.6	12.50	11.57	10.96
06 Sep 23	54001.6	12.54	11.51	10.74	08 Oct 19	54758.6	12.36	11.40	10.68
06 Sep 24	54002.6	12.63	11.57	10.78	08 Oct 21	54760.6	12.30	11.36	10.58
06 Sep 25	54003.6	12.59	11.55	10.78	08 Oct 23	54762.6	12.23	11.30	10.53
06 Sep 28	54006.6	12.62	11.60	10.85	08 Nov 02	54772.6	12.39	11.39	10.59
06 Sep 29	54007.6	12.67	11.63	10.89	08 Nov 03	54773.6	12.36	11.37	10.60
06 Sep 30	54008.6	12.66	11.65	10.92	08 Nov 09	54779.5	12.41	11.44	10.72
06 Oct 01	54009.6	12.65	11.60	10.91	08 Nov 11	54781.5	12.39	11.43	10.67
06 Oct 02	54010.6	12.63	11.61	10.91	08 Nov 17	54787.5	12.28	11.35	10.62
06 Nov 15	54054.5	12.66	11.67	10.98	08 Nov 20	54790.5	12.21	11.30	10.56
06 Nov 16	54055.6	12.69	11.69	11.01	09 Mar 01	54891.2	12.30	11.38	10.69
06 Nov 19	54058.5	12.66	11.64	10.98	09 Mar 13	54903.3	12.29	11.37	10.63
06 Nov 20	54059.5	12.67	11.65	10.97	09 Mar 17	54907.3	12.43	11.48	10.77
06 Nov 24	54063.5	12.57	11.54	10.88	11 Oct 27	55861.6	12.66	11.71	11.06
06 Nov 26	54065.5	12.74	11.72	11.04	11 Nov 01	55866.6	12.65	11.64	11.00
06 Nov 28	54067.6	12.60	11.61	10.92	11 Nov 09	55874.6	12.62	11.57	10.93
06 Nov 30	54069.5	12.61	11.62	10.94	11 Nov 14	55879.5	12.66	11.65	11.04
06 Dec 02	54071.5	12.69	11.67	10.97	11 Nov 19	55884.5	12.60	11.60	10.94
06 Dec 06	54075.5	12.69	11.69	11.00	13 Sep 26	56561.6	12.66	11.67	10.97
06 Dec 08	54077.4	12.63	11.64	10.93	13 Oct 14	56579.6	12.67	11.66	11.00
06 Dec 12	54081.5	12.70	11.68	11.01	13 Oct 25	56590.6	12.61	11.55	10.82
06 Dec 14	54083.5	12.60	11.59	10.92	13 Dec 18	56644.5	12.58	11.61	10.94
07 Feb 05	54137.3	12.69	11.72	11.08	13 Dec 23	56650.4	12.62	11.62	10.91
07 Mar 12	54172.3	12.73	11.77	11.15	14 Mar 11	56728.3	12.69	11.61	10.88
07 Oct 11	54385.6	12.60	11.59	10.90	14 Mar 13	56730.3	12.61	11.57	10.82
07 Oct 16	54390.6	12.61	11.60	10.90	14 Mar 18	56735.3	12.64	11.58	10.89
07 Oct 26	54400.6	12.57	11.54	10.79	14 Sep 30	56930.7	12.68	11.60	10.86
07 Nov 03	54408.6	12.44	11.44	10.71	14 Oct 18	56948.6	12.63	11.57	10.87

^aTypical errors of the near-IR magnitudes do not exceed 0.03 mag

Table 2. Detected emission lines, unreddened fluxes, and derived accretion rates assuming a visual extinction $A_V = 1$ mag.

Line ID	λ (μm)	Flux $\pm \Delta$ Flux (10^{-16} erg s $^{-1}$ cm $^{-2}$)	$\dot{M}_{acc}$ ($M_{\odot}$ yr $^{-1}$)
LBT/MODS			
H15	0.371	0.9 ± 0.3	–
H14	0.372	0.9 ± 0.3	–
H13	0.374	0.9 ± 0.3	–
H12	0.375	1.6 ± 0.3	–
H11	0.377	3.2 ± 0.4	–
H10	0.380	5.0 ± 0.4	5.0E-10
H9	0.384	8.4 ± 0.5	6.6E-10
H8	0.389	12.4 ± 0.5	7.1E-10
Ca II K	0.393	4.5 ± 0.2	2.1E-10
H7 + Ca II H	0.397	23.9 ± 0.5	–
He I	0.403	1.7 ± 0.3	1.0E-09
H δ	0.410	28.4 ± 0.4	1.2E-09
H γ	0.434	41.7 ± 0.4	1.3E-09
He I	0.447	4.4 ± 0.2	1.5E-09
Ti II	0.457	1.4 ± 0.3	–
He II	0.469	2.1 ± 0.3	–
He I	0.471	0.9 ± 0.2	1.8E-09
H β	0.486	66.6 ± 0.4	9.3E-10
He I	0.501	1.7 ± 0.2	9.1E-10
?	0.502	1.0 ± 0.2	–
?	0.517	1.8 ± 0.4	–
Mg I	0.518	0.8 ± 0.2	–
He I	0.588	9.5 ± 0.4	1.1E-09
[O I] ^a	0.630	3.2 ± 0.4	–
H α	0.656	567.2 ± 0.5	1.0E-09
He I	0.668	8.8 ± 0.8	2.4E-09
[S II] ^a	0.671	3.1 ± 0.8	–
[S II] ^a	0.673	3.4 ± 0.7	–
He I	0.707	6.7 ± 2.3	2.3E-09
He I	0.837	7.2 ± 1.2	–
O I	0.845	15.7 ± 2.7	9.9E-10
Ca II	0.850	22.5 ± 1.0	6.0E-10
Ca II	0.854	24.6 ± 0.7	6.0E-10
Ca II	0.866	20.7 ± 0.8	5.9E-10
He I	0.873	4.8 ± 1.4	–

Table 2—Continued

Line ID	λ (μm)	Flux $\pm \Delta$ Flux (10^{-16} erg s $^{-1}$ cm $^{-2}$)	$\dot{M}_{acc}$ ($M_{\odot}$ yr $^{-1}$)
Pa12	0.875	12.0 ± 2.2	–
He I	0.886	8.1 ± 1.1	–
Pa9	0.923	24.0 ± 8.5	1.4E-09
TNG/DOLORES			
He I	0.588	12.7 ± 1.5	1.5E-09
H α	0.656	656.8 ± 1.9	1.2E-09
He I	0.668	8.2 ± 1.5	2.2E-09
TNG/NICS			
Pa11	0.882	53.0 ± 11.0	–
Pa9	0.923	62.5 ± 10.0	–
Pa δ	1.004	85.1 ± 8.2	2.9E-09
He I	1.082	68.5 ± 6.3	8.3E-10
Pa γ	1.092	105.7 ± 10.0	2.5E-09
Pa β	1.281	262.3 ± 21.4	4.3E-09

Table 3. EW of the $H\beta$ emission line vs. B band underlying continuum measured in different epochs.

Source	Date (yy/mm/dd)	JD +2400000	EW ($\text{\AA}$)	B (mag)	Ref.
V1118 Ori	89 Jan 07	47533.5	- 13.6	14.6	1
	89 Jan 11	47537.5	- 13.8	14.0	1
	89 Feb 01	47558.5	- 9.7	13.7	1
	89 Dec 19	47879.5	- 7.0	13.6	1
	92 Dec 02	48958.5	- 28.7	15.6-16.2	2
	93 Feb 22	49040.5	- 21.4	15.6-16.2	2
	14 Mar 25	48958.5	- 36.0	18.4	3
EX Lup	93 Apr	...	- 24.0	13.2	4
	94 Mar 03	49414.5	- 9.4	12.0	4
	07 Jul	...	- 144	14.4	5

– Errors on EW and magnitude are typically $\sim 0.2 \text{ \AA}$ and $\lesssim 0.1$ mag, respectively

Note. — References to the Tables from 2 and 3: (1) Parsamian et al. 1996; (2) Parsamian et al. 2002; (3) this paper; (4) Herbig et al. 2001 (and reference therein); (5) Sipos et al. 2009; (6) Kun et al. 2011; (7) Aspin & Reipurth 2009; (8) Ojha et al. 2006; (9) Aspin et al. 2009; (10) Aspin et al. 2011b; (11) Aspin et al. 2011a; (12) Sicilia-Aguilar et al. 2008;

Table 4. EW of the H α emission line vs. R band underlying continuum measured in different epochs.

Source	Date (yy/mm/dd)	JD +2400000	H α EW ($\text{\AA}$)	B (mag)	Ref.
V1180 Cas	03 Feb 05	52675.5	- 300	16.7 ^a	6
	08 Aug 28	54706.5	- 900	17.5 ^a	6
	10 Dec 31	55561.5	- 530	17.0 ^a	6
V1647 Ori	04 Feb 14	53049.5	- 31.4	17.4 ^b	7
	04 Feb 22	53057.5	- 34.6	16.9	8
	04 Feb 23	53058.5	- 33.6	16.7	8
	04 Mar 10	53074.5	- 50.9	17.9 ^b	7
	04 Oct 06	53284.5	- 29.5	18.0 ^b	7
	04 Nov 16	53325.5	- 22.1	18.0 ^b	7
	04 Dec 12	53351.5	- 29.2	18.4 ^b	7
	05 Jan 08	53378.5	- 27.1	17.9 ^b	7
	05 Aug 30	53612.5	- 21.2	18.7 ^b	7
	05 Sep 25	53638.5	- 29.2	18.9 ^b	7
	05 Oct 13	53656.5	- 30.2	19.3 ^b	7
	05 Nov 18	53692.5	- 37.2	21.0 ^b	7
	05 Nov 27	53701.5	- 64.2	21.2 ^b	7
	05 Dec 25	53729.5	- 64.9	21.6 ^b	7
	06 Jan 05	53740.5	- 66.6	21.9 ^b	7
	06 Feb 16	53782.5	- 98.7	22.9 ^b	7
	08 Aug 31	54694.5	- 39	17.5 ^b	9
	11 Feb 02	55594.5	- 32.6	17.8 ^b	10
V2492 Cyg	10 Sep 25	55464.5	- 16.3	13.5	11
	10 Nov 05	55505.5	- 27.6	18.0	11
GM Cep	01 Jul 11	52101.5	- 10	12.7	12
	07 Apr 27	54217.5	- 19	13.5	12

– Notes and references are the same as in Table 3

^a Values of I_c magnitude estimated from a light-curve plot

^b Values of r' magnitude estimated from a light-curve plot

Table 5. Quiescence parameters of V1118 Ori.

Parameter		Value
Distance	d	400 pc
Spectral Type	SpT	M1e
Stellar radius	R_*	$1.29 R_\odot$
Stellar mass	M_*	$0.41 M_\odot$
Bolometric luminosity	L_{bol}	$2.1 L_\odot$
Visual extinction	A_V	1-2 mag
Mass accretion rate	$\dot{M}$	$1\text{-}3 \cdot 10^{-9} M_\odot \text{ yr}^{-1}$

Note. — The first four lines of the table list literature parameters, while the rest are derived in the present work.